


Clinical Profile When Testing for Cognition



Cognitive impairment may be caused by a number of underlying conditions.

Main Categories	Underlying conditions most commonly associated with cognitive impairment:	
Lifestyle	<ul style="list-style-type: none"> Poor diet Sedentarism Stress 	<ul style="list-style-type: none"> Unbalanced work-life style Substance abuse Polypharmacy
Cardiovascular	<ul style="list-style-type: none"> High blood pressure Heart failure COPD 	<ul style="list-style-type: none"> Arrhythmias Post-myocardial infarction Cardiorenal syndrome
Metabolic	<ul style="list-style-type: none"> Diabetes Vitamin deficiencies 	<ul style="list-style-type: none"> Dyslipidemia Liver diseases (cirrhosis, hepatitis, etc.)
Sleep Disorders	<ul style="list-style-type: none"> Chronic insomnia Sleep apnea 	<ul style="list-style-type: none"> Restless leg syndrome
Neurological Disorders	<ul style="list-style-type: none"> Chronic depression, anxiety Chronic pain TBI 	<ul style="list-style-type: none"> Multiple sclerosis Epilepsy Parkinson’s disease
Hormonal Imbalances	<ul style="list-style-type: none"> Hyper/hypothyroidism Menopause 	<ul style="list-style-type: none"> Post-stroke Dementias (vascular, Alzheimer’s disease, Frontotemporal, Lewy body)
Other	<ul style="list-style-type: none"> Post-surgery Post-chemo/radiation Vision and hearing impairment 	<ul style="list-style-type: none"> During hemodialysis Environmental toxins: (arsenic, lead, mercury, asbestos)